

Additive generators in interval-valued and intuitionistic fuzzy set theory

Glad Deschrijver

Fuzziness and Uncertainty Modeling Research Unit

Department of Mathematics and Computer Science, Ghent University

Krijgslaan 281 (S9), B-9000 Gent, Belgium

E-mail: Glad.Deschrijver@UGent.be

Homepage: <http://www.fuzzy.UGent.be>

Abstract

Intuitionistic fuzzy sets in the sense of Atanassov and interval-valued fuzzy sets can be seen as \mathcal{L} -fuzzy sets w.r.t. a special lattice \mathcal{L}^I . Deschrijver [2] introduced additive and multiplicative generators on \mathcal{L}^I based on a special kind of addition introduced in [3]. Actually, many other additions can be introduced. In this paper we investigate additive generators on \mathcal{L}^I as far as possible independently of the addition. For some special additions we investigate which t-norms can be generated by continuous additive generators which are a natural extension of an additive generator on the unit interval.

Keywords: intuitionistic fuzzy set, interval-valued fuzzy set, additive generator, addition on L^I , representable

1 Introduction

Triangular norms on $([0, 1], \leq)$ were introduced in [18] and play an important role in fuzzy set theory (see e.g. [12, 14] for more details). Generators are very useful in the construction of t-norms: any generator on $([0, 1], \leq)$ can be used to generate a t-norm. Generators play also an important role in the representation of continuous Archimedean t-norms on $([0, 1], \leq)$. Moreover, some properties of t-norms which have a generator can be related to properties of their generator. See e.g. [9, 13, 14, 15, 16] for more information about generators on the unit interval.

Interval-valued fuzzy set theory [11, 17] is an extension of fuzzy theory in which to each element of the universe a closed subinterval of the unit interval is assigned which approximates the unknown membership degree. Another extension of fuzzy set theory is intuitionistic fuzzy set theory introduced by Atanassov [1]. In [6] it is shown that intuitionistic fuzzy set theory is equivalent to interval-valued fuzzy set theory and that both are equivalent to L -fuzzy set theory in the sense of Goguen [10] w.r.t. a special lattice \mathcal{L}^I . In [2] additive and multiplicative generators on \mathcal{L}^I are investigated based on a special kind of addition introduced in [3]. In [8] another addition was defined. In fact, many more additions can be introduced. Therefore, in this paper we will investigate additive generators on \mathcal{L}^I as far as possible independently of the addition. For some special additions we will investigate which t-norms can be generated

by continuous additive generators which are a natural extension of an additive generator on the unit interval.

2 The lattice \mathcal{L}^I

Definition 2.1 We define $\mathcal{L}^I = (L^I, \leq_{L^I})$, where

$$L^I = \{[x_1, x_2] \mid (x_1, x_2) \in [0, 1]^2 \text{ and } x_1 \leq x_2\},$$

$$[x_1, x_2] \leq_{L^I} [y_1, y_2] \iff (x_1 \leq y_1 \text{ and } x_2 \leq y_2), \text{ for all } [x_1, x_2], [y_1, y_2] \text{ in } L^I.$$

Similarly as Lemma 2.1 in [6] it can be shown that \mathcal{L}^I is a complete lattice.

Definition 2.2 [11, 17] An interval-valued fuzzy set on U is a mapping $A : U \rightarrow L^I$.

Definition 2.3 [1] An intuitionistic fuzzy set on U is a set

$$A = \{(u, \mu_A(u), \nu_A(u)) \mid u \in U\},$$

where $\mu_A(u) \in [0, 1]$ denotes the membership degree and $\nu_A(u) \in [0, 1]$ the non-membership degree of u in A and where for all $u \in U$, $\mu_A(u) + \nu_A(u) \leq 1$.

An intuitionistic fuzzy set A on U can be represented by the \mathcal{L}^I -fuzzy set A given by

$$A : U \rightarrow L^I :$$

$$u \mapsto [\mu_A(u), 1 - \nu_A(u)],$$

In Figure 1 the set L^I is shown. Note that to each element $x = [x_1, x_2]$ of L^I corresponds a point $(x_1, x_2) \in \mathbb{R}^2$.




Figure 1: The grey area is L^I .

In the sequel, if $x \in L^I$, then we denote its bounds by x_1 and x_2 , i.e. $x = [x_1, x_2]$. The smallest and the largest element of \mathcal{L}^I are given by $0_{\mathcal{L}^I} = [0, 0]$ and $1_{\mathcal{L}^I} = [1, 1]$. We define

the relation \ll_{L^I} by $x \ll_{L^I} y \iff (x_1 < y_1 \text{ and } x_2 < y_2)$, for x, y in L^I . We define for further usage the sets

$$\begin{aligned} D &= \{[x_1, x_1] \mid x_1 \in [0, 1]\}; \\ \bar{L}^I &= \{[x_1, x_2] \mid (x_1, x_2) \in \mathbb{R}^2 \text{ and } x_1 \leq x_2\}; \\ \bar{D} &= \{[x_1, x_1] \mid x_1 \in \mathbb{R}\}; \\ \bar{L}_+^I &= \{[x_1, x_2] \mid (x_1, x_2) \in [0, +\infty[^2 \text{ and } x_1 \leq x_2\}; \\ \bar{D}_+ &= \{[x_1, x_1] \mid x_1 \in [0, +\infty[\}; \\ \bar{L}_{\infty,+}^I &= \{[x_1, x_2] \mid (x_1, x_2) \in [0, +\infty]^2 \text{ and } x_1 \leq x_2\}; \\ \bar{D}_{\infty,+} &= \{[x_1, x_1] \mid x_1 \in [0, +\infty]\}. \end{aligned}$$

Note that for any non-empty subset A of L^I it holds that

$$\begin{aligned} \sup A &= [\sup\{x_1 \mid x_1 \in [0, 1] \text{ and } (\exists x_2 \in [x_1, 1])([x_1, x_2] \in A)\}, \\ &\quad \sup\{x_2 \mid x_2 \in [0, 1] \text{ and } (\exists x_1 \in [0, x_2])([x_1, x_2] \in A)\}]. \end{aligned}$$

Definition 2.4 [4] *A t-norm on \mathcal{L}^I is a commutative, associative, increasing mapping $\mathcal{T} : (L^I)^2 \rightarrow L^I$ which satisfies $\mathcal{T}(1_{\mathcal{L}^I}, x) = x$, for all $x \in L^I$. A t-conorm on \mathcal{L}^I is a commutative, associative, increasing mapping $\mathcal{S} : (L^I)^2 \rightarrow L^I$ which satisfies $\mathcal{S}(0_{\mathcal{L}^I}, x) = x$, for all $x \in L^I$.*

In [4, 5, 7] the following classes of t-norms on \mathcal{L}^I are introduced: let T and T' be t-norms on $([0, 1], \leq)$, then the mappings $\mathcal{T}_{T,T'}$, \mathcal{T}_T , $\mathcal{T}_{T,t}$ and \mathcal{T}'_T given by, for all x, y in L^I ,

$$\begin{aligned} \mathcal{T}_{T,T'}(x, y) &= [T(x_1, y_1), T'(x_2, y_2)], \text{ (t-representable t-norms)} \\ \mathcal{T}_T(x, y) &= [T(x_1, y_1), \max(T(x_1, y_2), T(x_2, y_1))], \text{ (pseudo-t-representable t-norms)} \\ \mathcal{T}_{T,t}(x, y) &= [T(x_1, y_1), \max(T(t, T(x_2, y_2)), T(x_1, y_2), T(x_2, y_1))], \\ \mathcal{T}'_T(x, y) &= [\min(T(x_1, y_2), T(x_2, y_1)), T(x_2, y_2)], \end{aligned}$$

are t-norms on \mathcal{L}^I . The corresponding classes of t-conorms are given by, for all x, y in L^I ,

$$\begin{aligned} \mathcal{S}_{S,S'}(x, y) &= [S(x_1, y_1), S'(x_2, y_2)], \text{ (t-representable t-conorms)} \\ \mathcal{S}_S(x, y) &= [\min(S(x_1, y_2), S(x_2, y_1)), S(x_2, y_2)], \text{ (pseudo-t-representable t-conorms)} \\ \mathcal{S}_{S,t}(x, y) &= [\min(S(1-t, S(x_1, y_1)), S(x_1, y_2), S(x_2, y_1)), S(x_2, y_2)], \\ \mathcal{S}'_S(x, y) &= [S(x_1, y_1), \max(S(x_1, y_2), S(x_2, y_1))]. \end{aligned}$$

If for a mapping f on $[0, 1]$ and a mapping F on L^I it holds that $F(D) \subseteq \bar{D}$, and $F([a, a]) = [f(a), f(a)]$, for all $a \in L^I$, then we say that F is a natural extension of f to L^I . E.g. $\mathcal{T}_{T,T}$, \mathcal{T}_T , $\mathcal{T}_{T,t}$ and \mathcal{T}'_T are all natural extensions of T to L^I .

3 Additive generators on \mathcal{L}^I

In order to investigate additive generators on \mathcal{L}^I , a suitable addition on \bar{L}^I is needed. We assume from now on that $\oplus : (\bar{L}^I \cup \bar{L}_{\infty,+}^I)^2 \rightarrow \bar{L}^I$ satisfies the following natural properties, for all a, b in $\bar{L}^I \cup \bar{L}_{\infty,+}^I$,

- (i) \oplus is commutative,

- (ii) \oplus is associative,
- (iii) \oplus is increasing,
- (iv) $0_{\mathcal{L}^I} \oplus a = a$,

Note that from (iii) and (iv) it follows that $a \oplus b \geq_{L^I} a$, if $b \geq_{L^I} 0_{\mathcal{L}^I}$, for all a, b in $\bar{L}^I \cup \bar{L}_{\infty,+}^I$.

Definition 3.1 Let $\mathfrak{f} : L^I \rightarrow \bar{L}_{\infty,+}^I$ be a strictly decreasing function. The pseudo-inverse $\mathfrak{f}^{(-1)} : \bar{L}_{\infty,+}^I \rightarrow L^I$ of \mathfrak{f} is defined by, for all $y \in \bar{L}_{\infty,+}^I$,

$$\mathfrak{f}^{(-1)}(y) = \begin{cases} \sup\{x \mid x \in L^I \text{ and } \mathfrak{f}(x) \gg_{L^I} y\}, & \text{if } y \ll_{L^I} \mathfrak{f}(0_{\mathcal{L}^I}); \\ \sup(\{0_{\mathcal{L}^I}\} \cup \{x \mid x \in L^I \text{ and } (\mathfrak{f}(x))_1 > y_1 \\ \text{and } (\mathfrak{f}(x))_2 \geq (\mathfrak{f}(0_{\mathcal{L}^I}))_2\}), & \text{if } y_2 \geq (\mathfrak{f}(0_{\mathcal{L}^I}))_2; \\ \sup(\{0_{\mathcal{L}^I}\} \cup \{x \mid x \in L^I \text{ and } (\mathfrak{f}(x))_2 > y_2 \\ \text{and } (\mathfrak{f}(x))_1 \geq (\mathfrak{f}(0_{\mathcal{L}^I}))_1\}), & \text{if } y_1 \geq (\mathfrak{f}(0_{\mathcal{L}^I}))_1. \end{cases}$$

Definition 3.2 A mapping $\mathfrak{f} : L^I \rightarrow \bar{L}_{\infty,+}^I$ satisfying the following conditions:

- (AG.1) \mathfrak{f} is strictly decreasing;
- (AG.2) $\mathfrak{f}(1_{\mathcal{L}^I}) = 0_{\mathcal{L}^I}$;
- (AG.3) \mathfrak{f} is right-continuous in $0_{\mathcal{L}^I}$;
- (AG.4) $\mathfrak{f}(x) \oplus \mathfrak{f}(y) \in \mathcal{R}(\mathfrak{f})$, for all x, y in L^I , where

$$\begin{aligned} \mathcal{R}(\mathfrak{f}) = & \text{rng}(\mathfrak{f}) \cup \{x \mid x \in \bar{L}_{\infty,+}^I \text{ and } [x_1, (\mathfrak{f}(0_{\mathcal{L}^I}))_2] \in \text{rng}(\mathfrak{f}) \text{ and } x_2 \geq (\mathfrak{f}(0_{\mathcal{L}^I}))_2\} \\ & \cup \{x \mid x \in \bar{L}_{\infty,+}^I \text{ and } [(\mathfrak{f}(0_{\mathcal{L}^I}))_1, x_2] \in \text{rng}(\mathfrak{f}) \text{ and } x_1 \geq (\mathfrak{f}(0_{\mathcal{L}^I}))_1\} \\ & \cup \{x \mid x \in \bar{L}_{\infty,+}^I \text{ and } x \geq_{L^I} \mathfrak{f}(0_{\mathcal{L}^I})\}; \end{aligned}$$

- (AG.5) $\mathfrak{f}^{(-1)}(\mathfrak{f}(x)) = x$, for all $x \in L^I$;

is called an additive generator on \mathcal{L}^I .

Theorem 3.1 Let f be an additive generator on $([0, 1], \leq)$ and $\mathfrak{f} : L^I \rightarrow \bar{L}_{\infty,+}^I$ the mapping defined by, for all $x \in L^I$,

$$\mathfrak{f}(x) = [f(x_2), f(x_1)].$$

Then, for all $y \in \bar{L}_{\infty,+}^I$,

$$\mathfrak{f}^{(-1)}(y) = [f^{(-1)}(y_2), f^{(-1)}(y_1)]. \quad (1)$$

Lemma 3.2 Let $\mathfrak{f} : L^I \rightarrow \bar{L}_{\infty,+}^I$ be a mapping satisfying (AG.1), (AG.2), (AG.3) and (AG.5). Then, for all $x \in L^I$ such that $x_1 > 0$, it holds that $(\mathfrak{f}(x))_2 < (\mathfrak{f}(0_{\mathcal{L}^I}))_2$ and $(\mathfrak{f}(x))_1 < (\mathfrak{f}(0_{\mathcal{L}^I}))_1$.

Lemma 3.3 Let $\mathfrak{f} : L^I \rightarrow \bar{L}_{\infty,+}^I$ be a mapping satisfying (AG.1), (AG.2), (AG.3) and (AG.5). Then $(\mathfrak{f}([0, 1]))_1 = (\mathfrak{f}(0_{\mathcal{L}^I}))_1$ or $(\mathfrak{f}([0, 1]))_2 = (\mathfrak{f}(0_{\mathcal{L}^I}))_2$.

Corollary 3.4 Let $\mathfrak{f} : L^I \rightarrow \bar{L}_{\infty,+}^I$ be a mapping satisfying (AG.1), (AG.2), (AG.3), (AG.5) and $\mathfrak{f}(D) \subseteq \bar{D}_{\infty,+}$. Then $(\mathfrak{f}([0, 1]))_2 = (\mathfrak{f}(0_{\mathcal{L}^I}))_2$.

Theorem 3.5 Let $\mathfrak{f} : L^I \rightarrow \bar{L}_{\infty,+}^I$ be a continuous mapping satisfying (AG.1), (AG.2), (AG.3), (AG.5) and $\mathfrak{f}(D) \subseteq \bar{D}_{\infty,+}$. Then there exists an additive generator f on $([0, 1], \leq)$ such that, for all $a \in \bar{L}_{\infty,+}^I$,

$$\mathfrak{f}^{(-1)}(a) = [f^{(-1)}(a_2), f^{(-1)}(a_1)].$$

The following theorem can be shown independently of the addition \oplus used in (AG.4).

Theorem 3.6 A mapping $\mathfrak{f} : L^I \rightarrow \bar{L}_{\infty,+}^I$ is a continuous additive generator on \mathcal{L}^I such that $\mathfrak{f}(D) \subseteq \bar{D}_{\infty,+}$ if and only if there exists a continuous additive generator f on $([0, 1], \leq)$ such that, for all $a \in L^I$,

$$\mathfrak{f}(a) = [f(a_2), f(a_1)]. \quad (2)$$

4 Additive generators and t-norms on \mathcal{L}^I

We will give a sufficient condition for \oplus under which an additive generator associated to \oplus generates a t-norm. First we give a lemma.

Lemma 4.1 Let \mathfrak{f} be an additive generator on \mathcal{L}^I associated to \oplus . If, for all x, y, a in \bar{L}_+^I such that $x \leq_{L^I} a$ and $y \leq_{L^I} a \oplus a$,

$$y_2 \geq a_2 \implies \left((x \oplus y)_1 = (x \oplus [y_1, a_2])_1 \text{ or } \min((x \oplus y)_1, (x \oplus [y_1, a_2])_1) \geq a_1 \right) \quad (3)$$

and

$$y_1 \geq a_1 \implies \left((x \oplus y)_2 = (x \oplus [a_1, y_2])_2 \text{ or } \min((x \oplus y)_2, (x \oplus [a_1, y_2])_2) \geq a_2 \right) \quad (4)$$

then, for all $x \in L^I$ and $y \in \mathcal{R}(\mathfrak{f})$, we have that $\mathfrak{f}(x) \oplus \mathfrak{f}(\mathfrak{f}^{(-1)}(y)) \in \mathcal{R}(\mathfrak{f})$ and

$$\mathfrak{f}^{(-1)}(\mathfrak{f}(x) \oplus \mathfrak{f}(\mathfrak{f}^{(-1)}(y))) = \mathfrak{f}^{(-1)}(\mathfrak{f}(x) \oplus y).$$

Using Lemma 4.1, the following theorem can be shown.

Theorem 4.2 Let \mathfrak{f} be an additive generator on \mathcal{L}^I associated to \oplus . If (3) and (4) hold, for all x, y, a in L^I such that $x \leq_{L^I} a$ and $y \leq_{L^I} a \oplus a$, then the mapping $\mathcal{T} : (L^I)^2 \rightarrow L^I$ defined by, for all x, y in L^I ,

$$\mathcal{T}(x, y) = \mathfrak{f}^{(-1)}(\mathfrak{f}(x) \oplus \mathfrak{f}(y)),$$

is a t-norm on \mathcal{L}^I .

Theorem 4.3 Let f be an additive generator on $([0, 1], \leq)$. Then the mapping $\mathfrak{f} : L^I \rightarrow \bar{L}_{\infty,+}^I$ defined by, for all $x \in L^I$,

$$\mathfrak{f}(x) = [f(x_2), f(x_1)],$$

is an additive generator on \mathcal{L}^I associated to \oplus if and only if, for all x, y in L^I ,

$$\mathfrak{f}(x) \oplus \mathfrak{f}(y) \in (\text{rng}(f) \cup [f(0), +\infty])^2.$$

Theorem 3.6 and Theorem 3.1 show that no matter which operation \oplus is used in (AG.4), a continuous additive generator \mathfrak{f} on \mathcal{L}^I satisfying $\mathfrak{f}(D) \subseteq \bar{D}_{\infty,+}$ satisfies (2) and its pseudo-inverse satisfies (1). Therefore it depends on the operation \oplus which classes of t-norms on \mathcal{L}^I can have continuous additive generators that are a natural extension of additive generators on $([0, 1], \leq)$.

4.1 Additive generators based on $\oplus_{\mathcal{L}^I}$

Starting from the observation that the Łukasiewicz t-conorm S_W is given by $S_W(x, y) = \min(1, x + y)$, for all x, y in $[0, 1]$, and that the pseudo-t-representable t-conorm \mathcal{S}_{S_W} is given by $\mathcal{S}_{S_W}(x, y) = [\min(1, x_1 + y_2, x_2 + y_1), \min(1, x_2 + y_2)]$, for all x, y in L^I , the following definition of addition on \bar{L}^I is introduced in such a way that $\mathcal{S}_{S_W}(x, y) = \inf(1_{\mathcal{L}^I}, x \oplus_{\mathcal{L}^I} y)$, for all x, y in L^I .

Definition 4.1 [3] *We define the addition on $\bar{L}^I \cup \bar{L}_{\infty,+}^I$ by, for all x, y in $\bar{L}^I \cup \bar{L}_{\infty,+}^I$,*

$$x \oplus_{\mathcal{L}^I} y = [\min(x_1 + y_2, x_2 + y_1), x_2 + y_2],$$

where, for all $x \in \mathbb{R}$, $x + \infty = +\infty$ and $+\infty + \infty = +\infty$.

The other arithmetic operations introduced in [3] allow to write also some other important operations on \mathcal{L}^I , such as the Łukasiewicz t-norm, the product t-norm on \mathcal{L}^I and their residual implications, using a similar algebraic formula as their counterparts on $([0, 1], \leq)$.

Theorem 4.4 [2] *Let \mathfrak{f} be any generator on \mathcal{L}^I associated to $\oplus_{\mathcal{L}^I}$. Then the mapping $\mathcal{T} : (L^I)^2 \rightarrow L^I$ defined by, for all x, y in L^I ,*

$$\mathcal{T}(x, y) = \mathfrak{f}^{(-1)}(\mathfrak{f}(x) \oplus_{\mathcal{L}^I} \mathfrak{f}(y)),$$

is a t-norm on \mathcal{L}^I .

Theorem 4.5 [2] *Let \mathfrak{f} be a continuous additive generator on \mathcal{L}^I associated to $\oplus_{\mathcal{L}^I}$ for which $\mathfrak{f}(D) \subseteq \bar{D}_{\infty,+}$. Then there exists a t-norm T on $([0, 1], \leq)$ such that, for all x, y in L^I ,*

$$\mathfrak{f}^{(-1)}(\mathfrak{f}(x) \oplus_{\mathcal{L}^I} \mathfrak{f}(y)) = \mathcal{T}_T(x, y).$$

Thus, using $\oplus_{\mathcal{L}^I}$, only pseudo-t-representable t-norms on \mathcal{L}^I can have continuous additive generators \mathfrak{f} for which $\mathfrak{f}(D) \subseteq \bar{D}_{\infty,+}$. Other natural extensions of t-norms on $([0, 1], \leq)$ which have a continuous generator f cannot have a continuous additive generator on \mathcal{L}^I that is a natural extension of f .

4.2 Additive generators based on $\oplus_{\mathcal{L}^I}^t$

Now we discuss a second type of addition on \bar{L}^I which was introduced in [8]. Similarly as for $\oplus_{\mathcal{L}^I}$, we have that $\mathcal{S}_{S_W,t}(x, y) = \inf(1_{\mathcal{L}^I}, x \oplus_{\mathcal{L}^I}^t y)$, for all x, y in L^I .

Definition 4.2 [8] *Let $t \in [0, 1]$. Then we define the t-addition on $\bar{L}^I \cup \bar{L}_{\infty,+}^I$ by, for all x, y in $\bar{L}^I \cup \bar{L}_{\infty,+}^I$,*

$$x \oplus_{\mathcal{L}^I}^t y = [\min(1 - t + x_1 + y_1, x_1 + y_2, x_2 + y_1), x_2 + y_2].$$

Theorem 4.6 *Let $t \in [0, 1]$ and \mathfrak{f} be any generator on \mathcal{L}^I associated to $\oplus_{\mathcal{L}^I}^t$. Then the mapping $\mathcal{T} : (L^I)^2 \rightarrow L^I$ defined by, for all x, y in L^I ,*

$$\mathcal{T}(x, y) = \mathfrak{f}^{(-1)}(\mathfrak{f}(x) \oplus_{\mathcal{L}^I}^t \mathfrak{f}(y)),$$

is a t-norm on \mathcal{L}^I .

Theorem 4.7 *Let $t \in [0, 1]$ and \mathfrak{f} be a continuous additive generator on \mathcal{L}^I associated to $\oplus_{\mathcal{L}^I}^t$ for which $\mathfrak{f}(D) \subseteq \bar{D}_{\infty,+}$. Then there exists a t -norm T on $([0, 1], \leq)$ such that, for all x, y in L^I ,*

$$\mathfrak{f}^{(-1)}(\mathfrak{f}(x) \oplus_{\mathcal{L}^I}^t \mathfrak{f}(y)) = \mathcal{T}_{T, \mathfrak{f}^{-1}(1-t)}(x, y).$$

Similarly as for $\oplus_{\mathcal{L}^I}$, from Theorem 4.7 it follows that a t -norm \mathcal{T} on \mathcal{L}^I which is a natural extension of a t -norm on $([0, 1], \leq)$ generated by a continuous additive generator f can only have a continuous additive generator associated to $\oplus_{\mathcal{L}^I}^t$ which is a natural extension of f , if \mathcal{T} belongs to the class of t -norms $\mathcal{T}_{T,t}$.

4.3 Additive generators based on $\oplus'_{\mathcal{L}^I}$

Finally, we introduce the following addition on \bar{L}^I .

Definition 4.3 *We define the addition on $\bar{L}^I \cup \bar{L}_{\infty,+}^I$ by, for all x, y in $\bar{L}^I \cup \bar{L}_{\infty,+}^I$,*

$$x \oplus'_{\mathcal{L}^I} y = [x_1 + y_1, \max(x_1 + y_2, x_2 + y_1)].$$

This addition is closely related to the t -conorm \mathcal{S}'_{S_W} : for all x, y in L^I , $\mathcal{S}'_{S_W}(x, y) = \inf(1_{\mathcal{L}^I}, x \oplus'_{\mathcal{L}^I} y)$.

Theorem 4.8 *Let \mathfrak{f} be any generator on \mathcal{L}^I associated to $\oplus'_{\mathcal{L}^I}$ such that $\mathfrak{f}(0_{\mathcal{L}^I}) \in \bar{D}_{\infty,+}$. Then the mapping $\mathcal{T} : (L^I)^2 \rightarrow L^I$ defined by, for all x, y in L^I ,*

$$\mathcal{T}(x, y) = \mathfrak{f}^{(-1)}(\mathfrak{f}(x) \oplus'_{\mathcal{L}^I} \mathfrak{f}(y)),$$

is a t -norm on \mathcal{L}^I .

Theorem 4.9 *Let \mathfrak{f} be a continuous additive generator on \mathcal{L}^I associated to $\oplus'_{\mathcal{L}^I}$ for which $\mathfrak{f}(D) \subseteq \bar{D}_{\infty,+}$. Then there exists a t -norm T on $([0, 1], \leq)$ such that, for all x, y in L^I ,*

$$\mathfrak{f}^{(-1)}(\mathfrak{f}(x) \oplus'_{\mathcal{L}^I} \mathfrak{f}(y)) = \mathcal{T}'_T(x, y).$$

Similarly as for the two other additions, only t -norms on \mathcal{L}^I belonging to the class of t -norms \mathcal{T}'_T can have continuous additive generators \mathfrak{f} associated to $\oplus'_{\mathcal{L}^I}$ which are a natural extension of a continuous additive generator on $([0, 1], \leq)$.

5 Conclusion

In [3, 8] two kinds of arithmetic operations on \mathcal{L}^I are introduced. In [2] one of these kinds of operations is used to construct additive generators on \mathcal{L}^I . Since these are not the only possible ways to define addition, subtraction, multiplication and division on \mathcal{L}^I , we developed a new theory of additive generators on \mathcal{L}^I as much as possible independently of the addition needed. We found a sufficient condition for \oplus such that additive generators associated to \oplus generate t -norms on \mathcal{L}^I . We showed that continuous additive generators on \mathcal{L}^I which are a natural extension to \mathcal{L}^I of a generator on $([0, 1], \leq)$ can be represented in a unique way by the generator on $([0, 1], \leq)$. As a consequence, the choice of the operation \oplus determines which classes of t -norms on \mathcal{L}^I can have continuous additive generators which form a natural extension of a generator on the unit interval.

References

- [1] K. T. Atanassov, *Intuitionistic fuzzy sets*, Physica-Verlag, Heidelberg/New York, 1999.
- [2] G. Deschrijver, Additive and multiplicative generators in interval-valued fuzzy set theory, submitted.
- [3] G. Deschrijver, Algebraic operators in interval-valued fuzzy set theory, submitted.
- [4] G. Deschrijver, C. Cornelis and E. E. Kerre, On the representation of intuitionistic fuzzy t-norms and t-conorms, *IEEE Transactions on Fuzzy Systems*, **12**(1) (2004) 45–61.
- [5] G. Deschrijver and E. E. Kerre, Classes of intuitionistic fuzzy t-norms satisfying the residuation principle, *International Journal of Uncertainty, Fuzziness and Knowledge-Based Systems*, **11**(6) (2003) 691–709.
- [6] G. Deschrijver and E. E. Kerre, On the relationship between some extensions of fuzzy set theory, *Fuzzy Sets and Systems*, **133**(2) (2003) 227–235.
- [7] G. Deschrijver and E. E. Kerre, Implicators based on binary aggregation operators in interval-valued fuzzy set theory, *Fuzzy Sets and Systems*, **153**(2) (2005) 229–248.
- [8] G. Deschrijver and A. Vroman, Generalized arithmetic operations in interval-valued fuzzy set theory, *Journal of Intelligent and Fuzzy Systems*, in press.
- [9] W. M. Faucett, Compact semigroups irreducibly connected between two idempotents, *Proceedings of the American Mathematical Society*, **6** (1955) 741–747.
- [10] J. Goguen, L-fuzzy sets, *Journal of Mathematical Analysis and Applications*, **18**(1) (1967) 145–174.
- [11] M. B. Gorzalczyk, A method of inference in approximate reasoning based on interval-valued fuzzy sets, *Fuzzy Sets and Systems*, **21**(1) (1987) 1–17.
- [12] P. Hájek, *Metamathematics of fuzzy logic*, Kluwer Academic Publishers, Dordrecht, 1998.
- [13] E. P. Klement, R. Mesiar and E. Pap, Quasi- and pseudo-inverses of monotone functions, and the construction of t-norms, *Fuzzy Sets and Systems*, **104** (1999) 3–13.
- [14] E. P. Klement, R. Mesiar and E. Pap, *Triangular norms*, Kluwer Academic Publishers, Dordrecht, 2002.
- [15] C.-H. Ling, Representation of associative functions, *Publ. Math. Debrecen*, **12** (1965) 189–212.
- [16] P. S. Mostert and A. L. Shields, On the structure of semigroups on a compact manifold with boundary, *Annals of Mathematics*, **65** (1957) 117–143.
- [17] R. Sambuc, *Fonctions Φ -floues. Application à l'aide au diagnostic en pathologie thyroïdienne*, Ph. D. Thesis, Université de Marseille, France, 1975.
- [18] B. Schweizer and A. Sklar, *Probabilistic metric spaces*, North-Holland, New York, 1983.